

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed and, completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302 and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC, 20503				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE November 8, 1999		3. REPORT TYPE AND DATES COVERED Technical Report
4. TITLE AND SUBTITLE Metal Ion Complexation Studies of Novel 8-Hydroxyquinoline-Containing Diaza-18-Crown-6 Ligands and Analogues.			5. FUNDING NUMBERS N00014-98-1-0485 98PR05020-00	
6. AUTHOR(S) Jerald S. Bradshaw, Xian X. Zhang, Paul B. Savage, Krzysztof E. Krakowiak, and Reed M. Izatt				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry and Biochemistry Brigham Young University Provo, UT 84602			8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No 15	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Dr. Kelvin Higa (Program Director) Office of Naval Research 800 No. Quincy Street Arlington, VA 22217-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Interaction of some 8-hydroxyquinoline-substituted and related ligands (see Figure 1) with various metal ions was evaluated by a calorimetric titration technique at 25°C in MeOH. Bis-(8-hydroxyquinoline-2-ylmethyl)-substituted ligand 13 forms a very strong complex with Ba ²⁺ (log K = 11.6 in MeOH) and is highly selective for Ba ²⁺ over Na ⁺ , K ⁺ , Zn ²⁺ and Cu ²⁺ (selectivity factor > 10 ⁶). The ¹ H NMR spectral studies of the Ba ²⁺ complexes with bis-(8-hydroxyquinoline-2-ylmethyl)- and bis(5,7-dichloro-8-hydroxyquinoline-2-ylmethyl)-substituted diaza-18-crown-6 ligands (13 and 10) suggest that these complexes are cryptate-like structures with two overlapping hydroxyquinoline rings forming a pseudo second macroring.				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			16. PRICE CODE NA	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

OFFICE OF NAVAL RESEARCH

Grant N00014-98-1-0485

Technical Report No. 15

**Metal Ion Complexation Studies of Novel 8-Hydroxyquinoline-Containing
Diaza-18-Crown-6 Ligands and Analogues**

by

Jerald S. Bradshaw, Xian X. Zhang, Paul B. Savage,
Krzysztof E. Krakowiak, and Reed M. Izatt

Department of Chemistry and Biochemistry,
Brigham Young University, Provo, UT 84602

November 3, 1999

Reproduction in whole or in part is permitted for
any purpose of the University Government

This document has been approved for public release
and sale; its distribution is unlimited

Table 1. Log K , ΔH (kJ/mol), and $T\Delta S$ (kJ/mol) Values for Interactions of Macrocyclic Ligands with Metal Ions in Methanol Solution at 25.0 °C

ligand	cation	log K	ΔH	$T\Delta S$
1^a	Na ⁺	2.89	-14.1	2.4
	K ⁺	3.39	-24.4	-5.0
	Mg ²⁺	6.82	-2.5	36.4
	Ba ²⁺	3.60	-11.6	8.9
	Zn ²⁺	5.12	-11.4	-8.5
	Cu ²⁺	10.1	-92.5	-34.9
	Co ²⁺	5.14	-91.1	-61.8
2^a	Na ⁺	3.74	-26.4	-5.1
	K ⁺	6.61	-58.1	-20.4
	Mg ²⁺	b		
	Ca ²⁺	4.71	-25.2	1.7
	Sr ²⁺	4.67	-24.6	2.1
	Ba ²⁺	12.2	-76.1	-6.5
4	Mg ²⁺	5.7 ± 0.2	10.7 ± 0.9	43.2
	Co ²⁺	3.91 ± 0.08	-91.2 ± 0.5	-68.9
5	Mg ²⁺	5.02 ± 0.08	13.9 ± 0.9	42.6
	Co ²⁺	3.96 ± 0.06	-84.5 ± 0.5	-61.9
6	K ⁺	b		
	Ba ²⁺	3.57 ± 0.06	-29.4 ± 0.7	-9.0
	Zn ²⁺	4.80 ± 0.08	-64.8 ± 0.6	-37.4
	Cu ²⁺	5.01 ± 0.07	-57.6 ± 0.6	-29.0

...Continuation of Table 1

ligand	cation	log K	ΔH	T ΔS
8	Na ⁺	$\sim 3^c$	$\sim -5^c$	
	K ⁺	3.52 ± 0.03	-31.2 ± 0.4	-11.1
	Ba ²⁺	4.22 ± 0.05	-19.2 ± 0.8	4.9
	Zn ²⁺	> 5.5	-19.0 ± 0.5	> 12.4
	Cu ²⁺	4.28 ± 0.09	-55.3 ± 0.7	-30.9
13	Na ⁺	3.65 ± 0.01	-25.3 ± 0.2	-4.5
	K ⁺	5.88 ± 0.04^d	-55.6 ± 0.7^d	-22.0
	Ba ²⁺	11.6 ± 0.2^d	-73.0 ± 0.5^d	-6.8
	Zn ²⁺	4.92 ± 0.07^e	-95.7 ± 0.6	-67.6
	Cu ²⁺	4.39 ± 0.09	-100 ± 1	-74.9
14	Na ⁺	3.02 ± 0.05	-20.0 ± 0.6	-2.8
	K ⁺	3.82 ± 0.02	-47.8 ± 0.3	-26.0
	Ba ²⁺	4.87 ± 0.04	-26.4 ± 0.4	1.4
	Zn ²⁺	4.80 ± 0.08	-64.8 ± 0.6	-37.4
	Cu ²⁺	(Brown Precipitate)		


^a Bordunov, A. V.; Bradshaw, J. S.; Zhang, X. X.; Dalley, N. K.; Kou, X.-L.; Izatt, R. M. *Inorg. Chem.* **1996**, *35*, 7229.

^bNo measurable heat other than heat of dilution indicating that ΔH or/and log K is small.

^cEstimated by a competitive calorimetric titration with Zn²⁺.

^dDetermined by a competitive calorimetric titration.

^eWhen [Zn²⁺]/13 \geq 2, a white precipitate formed.




1 X = Cl, Y = H

4 X = Y = H


5 X = CH₃, Y = H

6 X = H, Y = OH




2 X = Cl

13 X = H



8



14

Figure 1. 5-Chloro-8-hydroxyquinoline(CHQ)-substituted Diaza-18-crown-6 Ligand Analogues of **1** and **2** Used in this Study